

Appendix 2. Seasonal changes in the utilisation of the nine vegetation types for grazing (cf. Table 2; type 9 = *Schoenetum*). For each observation date, the time spent grazing in each vegetation type was expressed as percentage of all grazing observations (see Appendix 1 for calculations).

Date	Percentage of observations in each vegetation type								
	1	2	3	4	5	6	7	8	9
05.06.2002	2.9	18.4	31.1	38.8	1.0	0.0	0.0	7.9	0.0
07.06.2003	0.0	0.0	0.0	37.2	21.1	0.0	0.7	24.9	16.1
12.06.2002	2.2	5.2	16.8	16.8	27.8	5.2	0.7	24.7	0.5
20.06.2002	0.1	1.6	5.5	12.6	46.7	14.0	0.0	19.5	0.0
26.06.2002	0.0	7.0	16.7	12.7	39.4	0.3	18.0	5.9	0.0
05.07.2002	0.9	10.5	22.9	8.5	20.7	5.0	2.6	23.6	5.4
12.07.2002	7.7	5.2	12.0	0.2	10.8	44.8	0.9	18.4	0.2
17.07.2002	3.7	6.9	27.3	4.2	9.7	9.9	0.1	36.8	1.5
24.07.2002	6.3	7.2	27.2	3.4	18.8	9.7	0.3	27.0	0.1
25.07.2002	0.1	0.0	15.7	6.3	1.6	0.0	1.3	57.1	17.9
31.07.2002	6.1	5.1	37.6	1.8	0.0	0.0	0.5	38.7	10.1
09.08.2002	7.7	9.2	22.5	4.2	23.8	0.0	4.1	26.7	1.7
14.08.2002	25.5	11.4	48.5	0.7	0.0	0.0	0.3	4.3	9.3
23.08.2002	5.9	11.1	39.9	12.3	9.3	0.0	1.2	17.2	3.1
30.08.2002	2.6	6.6	43.4	0.3	14.2	5.7	7.8	16.0	3.4